



Limb Asymmetry Characteristics Amongst Men's and Women's NCAA Division II Soccer Athletes

Gavin Scott¹ and Brett Crossland^{2*}

¹Assistant Mens Soccer Coach, Young Harris College, 1 College St Young Harris, GA 30582, United States.

^{2*}Assistant Professor, Department of Athletic Training & Exercise Physiology, Midwestern State University, 3410 Taft Blvd. Wichita Falls, TX 76308, United States.

Article Details

Article Type: Research Article

Received date: 30th May, 2025

Accepted date: 01st July, 2025

Published date: 03rd July, 2025

***Corresponding Author:** Brett Crossland, Assistant Professor, Department of Athletic Training & Exercise Physiology, Midwestern State University, 3410 Taft Blvd. Wichita Falls, TX 76308, United States.

Citation: Scott, G., & Crossland, B., (2025). Limb Asymmetry Characteristics Amongst Men's and Women's NCAA Division II Soccer Athletes. *J Rehab Pract Res*, 6(2):178. <https://doi.org/10.33790/jrpr1100178>

Copyright: ©2025, This is an open-access article distributed under the terms of the [Creative Commons Attribution License 4.0](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Differences in the performance and function of one limb with respect to the other is known as interlimb asymmetry, while intralimb asymmetry deals with differences in muscle forces within the same limb. Both forms of asymmetry are common in the sport of soccer due to its distinct characteristics. These differences in strength between dominant and nondominant limbs or muscle groups may lead to increased injury rates in soccer athletes. The purpose of this study was to compare the magnitude of interlimb and intralimb asymmetry between NCAA Division II men's and women's soccer players. Thirty participants (male = 15, female = 15) completed a testing protocol in which concentric knee extensor and flexor peak torque was measured. Three isometric contractions lasting five seconds each were completed by all participants for each action on the dominant and nondominant limb. Interlimb asymmetries were observed in each sex (Male: Quad (Q) = $10.62 \pm 8.89\%$, Hamstring (H) = $13.33 \pm 8.43\%$. Females: Q = $7.70 \pm 7.63\%$ H = $8.56 \pm 6.04\%$) in both the Q and H muscle groups with the dominant limb being stronger. When comparing the sexes, males possessed significantly larger ($p = 0.04$) interlimb hamstring imbalances as compared to females. While females observed slightly elevated intralimb (H:Q) ratios for both limbs as compared to males, no significant differences were observed in H:Q ratio between the groups in the dominant ($p = 0.485$) or nondominant leg ($p = 0.195$). Results of interest indicate that male participants possessed significantly larger magnitude of interlimb imbalance as compared to females, and males may observe a greater benefit from focused training aimed at reducing interlimb asymmetries specifically in the hamstring group. This research only investigated isometric contractions which may pose as a limitation to the inference of results.

Key words: Strength Imbalance, Injury Rate, Soccer Performance, Football

Introduction

Differences in the performance and function of one limb with respect to the other is known as interlimb asymmetry, while intralimb

asymmetry involves differences in muscle forces within the same limb [1]. It is not uncommon for athletes participating in sport to show varying degrees of asymmetry. This is often observed in the sport of soccer with its distinct physical, technical, positional, and limb favouring tendencies which can lead to higher magnitudes of asymmetry compared to other sports [2]. The unilateral movements commonly utilized in soccer such as kicking, jumping, and changing direction may lead to unique loading patterns that can result in increased inter and intralimb asymmetry as compared to other team sports [3]. Previous research has investigated limb asymmetry and the potential effects it may have on injury risk and performance [4-7].

In soccer, muscular strength of the knee joint is vital in supporting the individual's body weight and center of mass while performing dynamic movements [8]. With regard to intralimb asymmetry in soccer, particular attention has been paid to hamstring and quadriceps discrepancies (H:Q ratio) due to the hypothesized relationship with injury risk [4, 9, 10]. Typical H:Q ratios in athletes can range from 0.5 to 0.8 with H:Q ratios less than 0.6 potentially leading to an increased risk of anterior cruciate ligament (ACL) and hamstring injury in participants [11-13]. Interlimb, or bilateral H:Q ratio differences, have been investigated with some research finding differences in field athletes when compared to court athletes with others finding no differences between the sexes [14, 15].

Interlimb asymmetry is not uncommon in humans and can occur across a number of factors such as strength, flexibility, balance, and power [1]. These asymmetries, typically exacerbated by overuse, can be caused by injury, dominant limb choice, playing position, or anthropometric imbalances [16]. While easy to identify, understanding the associated consequences of such asymmetries is challenging due to inconsistent findings within the research [5]. With regard to knee injuries, research has clearly shown females to be at a higher risk when compared to males, yet understanding the mechanisms that lead to this discrepancy still remain unclear [17]. Intralimb imbalances between the hamstring and quadriceps muscle groups are hypothesized to play a major role in the increased risk of

knee injury in females but this is likely the result of multiple factors such as hormonal fluctuations, neuromuscular differences, developmental changes, and anatomical differences [18]. While H:Q ratio is commonly accepted as a risk factor, recent research suggests H:Q ratio may not to be an independent risk factor or ACL injury further confusing researchers and practitioners [19].

Needless to say, research outcomes specific to muscular imbalances and athletic performance are varied, with no clear guidelines for practitioners or athletes to follow. The goal of this research project is to directly compare intra and interlimb imbalances between male and female athletes of similar age and experience level. In doing so, the researchers hope to increase the body of knowledge on muscular imbalances and help coaches and athletes better optimize training strategies. Recent data suggests the physiological demands between men's and women's soccer differ as female athletes typically cover a shorter total distance but at higher relative intensities when compared to males as measured by GPS data [20, 21]. Women's soccer is one of the fastest growing sports in the world, therefore determining potential differences in limb asymmetry between sexes may provide valuable insight regarding training methodological differences between the sexes [5, 22, 23].

Further research could provide insight for future programming of injury prevention protocols, rehabilitation techniques, strength and conditioning plans, and planning of training sessions. Therefore, the purpose of this study is to determine the magnitude of limb asymmetry between National Collegiate Athletic Association (NCAA) Division II collegiate men's and women's soccer players. The increasing popularity of women's soccer and the conflicting research regarding sex differences in asymmetry magnitude provides the basis for the need of this research study. Furthermore, these results will contribute to the body of research that currently exists while providing relevant information to players, coaches, trainers, and other healthcare professionals regarding potential differences between male and female soccer players.

Methodology

To examine sex differences in isometric limb asymmetry a quantitative comparative design was utilized. Participants were measured for isometric knee extensor and flexor strength to determine potential imbalances by performing three maximal isometric knee extension and flexion contractions for each leg. Participants recruited for this research project were male and female NCAA Division II soccer players. All research procedures were approved by the Institutional Review Board at Midwestern State University prior to participant recruitment.

Participants

Thirty participants ($n = 30$) including fifteen males ($n = 15$) and fifteen females ($n = 15$) were recruited using convenience sampling. Participants were screened to include those who: 1) were current NCAA Division II soccer players, 2) identify as an individual primarily playing any position other than goalie, 3) were 18-24 years of age, 4) were deemed healthy as determined by a medical questionnaire and PAR Q+ questionnaire, and 5) had a minimum of 5 years' experience playing competitive soccer. Participants were also screened to include those without: 1) lower-limb muscle or joint injury in past year that has resulted in more than five missed training sessions and 2) musculoskeletal injuries that limit the ability to perform the isometric strength testing. All participants were instructed that they were free to remove themselves from the study at any time.

Study Protocol

A single laboratory visit was utilized to complete all testing protocols. All testing was completed in the morning following 24 hours with no soccer training prior to eating or drinking. Height (cm), mass (kg) and body composition (BF%) were collected using a Seca mBCA 554 body composition analyzer (GmbH, Hamburg, Germany) which has been validated as an accurate and reliable method of

measuring body composition via bioelectrical impedance analysis [24]. Participants completed a 5-minute warm-up at a self-selected cadence on an electronically braked cycle ergometer (Ergomedic 828E; Monark Exercise AB, Vansbro, Sweden). Following the warm-up protocol, participants completed isometric strength testing on an isokinetic dynamometer (Biodex Medical Systems, Inc., Shirley NY, USA). Three maximal isometric contractions were performed for knee extension and flexion on each leg. Each isometric contraction was performed for 5 seconds with one minute of rest given between repetitions. The determination of testing order for limb and joint action performed first was at random. For all knee extension measurements seat orientation was 90°, seatback tilt was 80°, and knee angle was set at 90° of knee flexion with the lateral femoral epicondyle aligned with the axis of the dynamometer. For all knee flexion measurements seat orientation was 90°, seatback tilt was 80°, and knee angle was set at 60° of knee flexion with the lateral femoral condyle aligned with the axis of the dynamometer. Participants were given the opportunity to familiarize themselves with all testing protocols prior to data collection. Peak torque ($N \cdot m$) exerted for each repetition on dominant and nondominant legs for flexion and extension was recorded with the highest score on each limb and joint action being included for analysis.

Statistical Analysis

Statistical software (IBM SPSS Statistics v.24, Armonk, NY) was used to analyze all asymmetry measures between the sexes. Interlimb asymmetry was calculated using the following equation adapted by Impellizzeri et al. [25]: $\text{Asymmetry} = (\text{stronger} - \text{weaker}) / \text{stronger} * 100$. Intralimb (H:Q) asymmetry was calculated by dividing peak torque achieved by the hamstrings by peak torque achieved by the quadriceps. A one-way analysis of variance (ANOVA) was used to determine if statistical differences exist between groups for intra and interlimb asymmetry. The significance level was set *a priori* at 0.05.

Results

Fifteen male and fifteen female NCAA Division II soccer players completed the IRB approved testing protocol with no adverse events recorded. All participants reported as being right leg dominant, therefore no differentiation between dominant and non-dominant limb was required for reporting purposes. Descriptive characteristics of male participants ($n = 15$) and female ($n = 15$) are noted in Table 1.

Characteristics	Male	Female
Age (yrs)	21.8 ± 1.3	19.3 ± 1.0
Height (cm)	178.2 ± 7.6	164.8 ± 4.8
Weight (kg)	76.5 ± 8.4	64.8 ± 6.6
Body Composition (BF%)	16.0 ± 3.2	26.9 ± 4.3
Competitive Soccer Experience (yrs)	16.3 ± 2.3	13.9 ± 1.8

Table 1: Descriptive Characteristics of Male and Female Participants

Values are presented as mean ± SD; yrs = years, BF % = percentage of body fat.

Interlimb Asymmetry

Values used in the comparative analysis for asymmetry of the quadricep and hamstring muscles were derived from the equation: $\text{asymmetry} = (\text{stronger limb} - \text{weaker limb}) / \text{stronger limb} * 100$ [25]. While not statistically different ($p = 0.34$), interlimb asymmetries for the quadriceps muscle group were elevated in the male participants. Male participants possessed an average quadricep interlimb asymmetry magnitude of $10.62\% \pm 5.99$ compared to $7.70\% \pm 4.63$ for female participants.

With regards to hamstring interlimb asymmetry, male participants reported a significantly higher ($p = 0.04$) asymmetry compared to the

female participants ($13.33\% \pm 5.45$ vs $8.56\% \pm 4.74$). These results indicate higher discrepancies of strength in the hamstring muscle group between the dominant and nondominant limbs in males.

Values for interlimb imbalances for male and female participants can be seen in Figure 1.




Figure 1: Interlimb Asymmetry Magnitudes

Note. Interlimb asymmetry values for male and female hamstring and quadricep muscle groups. Hamstring interlimb asymmetry was significantly lower ($p = 0.04$) in the female participants. * Denotes significantly different than Male values.

Intralimb Asymmetry

H:Q ratio asymmetry was calculated by dividing peak torque achieved by the hamstrings by peak torque achieved by the quadriceps. While females showed elevated H:Q ratios for both

limbs as compared to males, no significant differences were observed in H:Q ratio between the sexes in the dominant ($p = 0.485$) or nondominant legs ($p = 0.195$). Values for H:Q ratios between sexes for can be found in Figure 2.




Figure 2: H:Q Ratio Amongst Participants

Note. H:Q ratios for male and female participants. No significant differences were observed.

Discussion

The purpose of this study was to investigate the magnitude of intra and interlimb asymmetry between NCAA Division II collegiate men's and women's soccer players. The key findings of this investigation were: 1) a significantly higher interlimb hamstring asymmetry in male participants as compared to the female participants, and 2) no significant differences in H:Q ratio between male and female participants. Amongst the measured participants, it appears that the male participants exhibited significantly higher levels of both interlimb asymmetry as compared to female participants, and practically significant higher H:Q ratio.

Interlimb asymmetry is commonly referred to as the differences in the performance and function of one limb or muscle group with respect to the other [1]. It is not uncommon for soccer athletes to exhibit varying degrees of interlimb asymmetry as a result of the repetitive movement patterns performed while playing the sport [2, 3]. These imbalances may have a negative effect on performance and increase an individual's risk of experiencing a lower-body injury [26, 27]. Researchers have observed a positive correlation between interlimb strength asymmetries and decreased sprint time as well as change of direction time in elite under 21 male and female soccer players [26]. Izovska et al. [27] observed that independent of sex, soccer players

possessing more than a 10% bilateral knee extensor asymmetry experienced higher rates of lower-body injuries. Based on this information, it appears that the male participants who participated in the current study would be at higher risk of experiencing negative outcomes such as decreased performance and increased injury risk as compared to the female participants.

Carey et al. [28] observed that elite lever soccer athletes complete approximately 83% of their touches on the ball with their dominant limb. This unbalanced preference of one limb over the other is likely one of the determining factors in the magnitude of imbalance observed in soccer athletes [29]. For both the male and female participants who took part in the study, the dominant limb produced higher levels of torque for all measured variables as compared to the nondominant limb. While common, these imbalances do seem to pose an injury risk as multiple researchers have observed increased injury risk in the dominant limb in soccer athletes [29-31]. Given this information, it appears that interlimb imbalances should be an area of focus for strength and conditioning and sports medicine staff as it relates to potentially reducing the risk of injury in soccer athletes.

One concerning finding in this project is both the men and women measured in the current study fell below a H:Q ratio of 0.60 which has been hypothesised to lead to reduced performance and increased risk of injury [13]. While not uncommon amongst soccer athletes, this does warrant attention. Research suggests that female athletes may be up to 8 times more likely to experience and ACL tear as compared to their male counterparts [32]. Females typically possess greater imbalances between hamstring and quadricep strength as compared to males, and this increased imbalance has been hypothesized to play a large role in the increased incidence of ACL injuries observed in females although this is highly debated [33, 34]. In a review by Hewett et al. [33], researchers observed females H:Q ratio was 8.8% lower when compared to males across all velocities indicating reduced hamstring strength as compared to the quadriceps. Females in the current study measured higher for both the dominant and nondominant limbs for H:Q ratio indicating less intralimb asymmetry as compared to the male participants. Although not statistically significant, these results provide some practical significance in that they differ from previous research in which it has been established that males typically possess a higher H:Q ratio as compared to females [35-37]. Given the known injury discrepancies between male and female participants and a growing knowledge of injury prevention techniques as it relates to the knee joint, one likely explanation for higher H:Q ratios observed in females may be an increased focus on preventative training. Participants in the current study averaged more than 15 years of experience playing competitive soccer and have also had varied access to structured strength and conditioning programs over this time frame.

Isokinetic and isometric methodologies are frequently used in measuring both inter and intralimb imbalances. While some research suggests sex differences between the varying speeds of contraction, the use of only isometric measures alone is a potential limitation of the current research study [33]. For the purposes of this investigation researchers were interested in maximal strength as opposed to dynamic strength, therefore, isometric contractions were chosen for testing over isokinetic. The use of NCAA Division II soccer athletes as participants limits the scope of the findings to individuals of similar ages and with similar training backgrounds. Participants in the current research study had over 15 years of experience participating in competitive soccer. Given the repeated movement patterns that are typically performed within the sport of soccer in addition to individuals preferring to use their dominant limb, the imbalances observed can only be conferred to individuals possessing a similar training background in soccer although some recent research suggests imbalance magnitude is not affected by training ages in male soccer athletes [38]. Due to the specific age and training inclusion criteria the sample size for this project is smaller

than desired therefore the results should be interpreted with caution, and may be only applicable to individuals with similar characteristics.

Conclusion

While research regarding muscular imbalance and negative performance or injury outcomes is mixed, there is little debate that a minimizing the magnitude of imbalances should be a focus of any training regime. The current research findings observed varying levels of intra and interlimb imbalances in both male and female Division II soccer athletes. The magnitude of imbalances an individual may experience depends on a number of factors including playing position, playing style, and training age [3]. It is the recommendation of the researchers that coaches and training staff should aim to identify imbalances and track those variables over time in order to determine if an individual's current training practices are increasing or decreasing any observed imbalances. While intra and interlimb imbalances appear to be a consequence of repetitive movements thus a common occurrence in soccer, minimizing these imbalances should be a priority in training.

Future research may focus on tracking imbalances over extended periods of time in order to begin to better understand additional factors that may lead to increases in measured differences. Factors such as weekly training load, number of matches played, and playing surface type could provide more insight on the synergistic effect of all training variables. In addition, specific rehabilitation practices aimed at reducing imbalances should be investigated to determine optimal methods of reducing or preventing both inter and intralimb imbalances in beginner and experienced athletes alike.

Competing Interests: The authors declare that they have no competing interests.

References

1. Bishop, C., Read, P., Chavda, S., & Turner, A. (2016). Asymmetries of the lower limb: the calculation conundrum in strength training and conditioning. *Strength and Conditioning Journal*, 38(6), 27–32. <https://doi.org/10.1519/ssc.0000000000000264>
2. Read, P. J., Oliver, J. L., De Ste Croix, M. B. A., Myer, G. D., & Lloyd, R. S. (2018). A prospective investigation to evaluate risk factors for lower extremity injury risk in male youth soccer players. *Scandinavian Journal of Medicine and Science in Sports*, 28(3), 1244–1251. <https://doi.org/10.1111/sms.13013>
3. Nicholson, G., Bennett, T., Thomas, A., Pollitt, L., Hopkinson, M., Crespo, R., Robinson, T., & Price, R. J. (2022). Inter-limb asymmetries and kicking limb preference in English premier league soccer players. *Frontiers in Sports and Active Living*, 4. <https://doi.org/10.3389/fspor.2022.982796>
4. Bishop, C., Coratella, G., & Beato, M. (2021). Intra- and inter-limb strength asymmetry in soccer: A comparison of professional and under-18 players. *Sports*, 9(9), 129. <https://doi.org/10.3390/sports9090129>
5. Guan, Y., Bredin, S. S. D., Taunton, J., Jiang, Q., Wu, N., & Warburton, D. E. R. (2022). Association between inter-limb symmetries in lower-limb functional performance and sport injury: A systematic review of prospective cohort studies. *Journal of clinical medicine*, 11(2), 360. <https://doi.org/10.3390/jcm11020360>
6. Jaiyesimi, A. O., & Jegede, J. A. (2005). Hamstring and quadriceps ratio: effect of age and gender. *Journal of the Nigeria Society of Physiotherapy*, 15(2), 54–58. <https://academicjournals.org/journal/JNSP/how-to-cite-article/93E96E765319>
7. Pardos-Mainer, E., Bishop, C., Gonzalo-Skok, O., Nobari, H., Pérez-Gómez, J., & Lozano, D. (2021). Associations between inter-limb asymmetries in jump and change of direction speed tests and physical performance in adolescent female soccer players. *International Journal of Environmental Research and Public Health*, 18(7), 3474. <https://doi.org/10.3390/ijerph18073474>

8. Myer, G. D., Ford, K. R., & Hewett, T. E. (2004). Rationale and clinical techniques for anterior cruciate ligament injury prevention among female athletes. *PubMed*, 39(4), 352–364. <https://pubmed.ncbi.nlm.nih.gov/15592608>
9. Beato, M., Young, D., Stiff, A., & Coratella, G. (2021). Lower-limb muscle strength, anterior-posterior and inter-limb asymmetry in professional, elite academy and amateur soccer players. *Journal of Human Kinetics*, 77, 135–146. <https://doi.org/10.2478/hukin-2020-0058>
10. Jeon, K., Chun, S., & Seo, B. (2016). Effects of muscle strength asymmetry between left and right on isokinetic strength of the knee and ankle joints depending on athletic performance level. *Journal of Physical Therapy Science*, 28(4), 1289–1293. <https://doi.org/10.1589/jpts.28.1289>
11. Dauty, M. (2016). Prediction of hamstring injury in professional soccer players by isokinetic measurements. *Muscles Ligaments and Tendons Journal*. <https://doi.org/10.11138/mltj/2016.6.1.116>
12. Sangnier, S., & Tourny-Chollet, C. (2007). Comparison of the decrease in strength between hamstrings and quadriceps during isokinetic fatigue testing in semiprofessional soccer players. *International Journal of Sports Medicine*, 28(11), 952–957. <https://doi.org/10.1055/s-2007-964981>
13. Yıldız, M., & Kale, M. (2017). The effects of kicking leg preference on the bilateral leg strength asymmetries of amateur football players. *Isokinetics and Exercise Science*, 26(1), 37–42. <https://doi.org/10.3233/ies-171159>
14. Cheung, R., Smith, A., & Wong, D. (2012). H:Q ratios and bilateral leg strength in college field and court sports players. *Journal of Human Kinetics*, 33(2012), 63–71. <https://doi.org/10.2478/v10078-012-0045-1>
15. Yilmaz, E., Aydin, T., Kilic, S., & Toluk, Ö. (2023). Bilateral knee joint isokinetic muscle strength and angle specific balance ratio in soccer players. *Deutsche Zeitschrift für Sportmedizin*, 74(7), pp. 234–241. <https://doi.org/10.5960/dzsm.2023.580>
16. Heil, J., Loffing, F., & Büsch, D. (2020). The influence of exercise-induced fatigue on inter-limb asymmetries: A systematic review. *Sports Medicine*, 6(1), 39. <https://doi.org/10.1186/s40798-020-00270-x>
17. Zech, A., Hollander, K., Junge, A., Steib, S., Groll, A., Heiner, J., Nowak, F., Pfeiffer, D., & Rahlf, A. L. (2022). Sex differences in injury rates in team-sport athletes: A systematic review and meta-regression analysis. *Journal of Sport and Health Science*, 11(1), 104–114. <https://doi.org/10.1016/j.jshs.2021.04.003>
18. Mancino, F., Kayani, B., Gabr, A., Fontalis, A., Plastow, R., & Haddad, F. S. (2024). Anterior cruciate ligament injuries in female athletes: risk factors and strategies for prevention. *Bone & Joint Open*, 5(2), 94–100. <https://doi.org/10.1302/2633-1462.52.BJO-2023-0166>
19. Kellis, E., Sahinis, C., & Baltzopoulos, V. (2023). Is hamstrings-to-quadriceps torque ratio useful for predicting anterior cruciate ligament and hamstring injuries? A systematic and critical review. *Journal of Sport and Health Science*, 12(3), 343–358. <https://doi.org/10.1016/j.jshs.2022.01.002>
20. Jagim, A. R., Murphy, J., Schaefer, A. Q., Askow, A. T., Luedke, J. A., Erickson, J. L., & Jones, M. T. (2020). Match demands of women's collegiate soccer. *Sports*, 8(6), 87. <https://doi.org/10.3390/sports8060087>
21. Stølen, T., Chamari, K., Castagna, C., & Wisløff, U. (2005). Physiology of soccer: An update. *Sports Medicine (Auckland, N.Z.)*, 35(6), 501–536. <https://doi.org/10.2165/00007256-200535060-00004>
22. Crossley, K.M., Patterson, B.E., Culvenor, A.G., Bruder, A.M., Mosler, A.B., & Mentiplay, B.F. (2020). Making football safer for women: A systematic review and meta-analysis of injury prevention programmes in 11 773 female football (soccer) players. *British Journal of Sports Medicine*, 54, 1089–1098. <http://doi.org/10.1136/bjsports-2019-101587>
23. Kong, P. W., & Burns, S. F. (2009). Bilateral difference in hamstrings to quadriceps ratio in healthy males and females. *Physical Therapy in Sport*, 11(1), 12–17. <https://doi.org/10.1016/j.ptsp.2009.09.004>
24. Bosy-Westphal, A., Jensen, B., Braun, W., Pourhassan, M., Gallagher, D., & Müller, M. J. (2017). Quantification of whole-body and segmental skeletal muscle mass using phase-sensitive 8-electrode medical bioelectrical impedance devices. *European Journal of Clinical Nutrition*, 71(9), 1061–1067. <https://doi.org/10.1038/ejcn.2017.27>
25. Impellizzeri, F.M., Rampinini, E., Castagna, C., Martino, F., Fiorini, S., Wisloff, U., (2008). Effect of plyometric training on sand versus grass on muscle soreness, jumping and sprinting ability in soccer players. *Br J Sports Med*. 42(1).doi:10.1136/bjism.2007.038497
26. Coratella, G., Beato, M., & Schena, F. (2018). Correlation between quadriceps and hamstrings inter-limb strength asymmetry with change of direction and sprint in U21 elite soccer-players. *Human Movement Science*, 59, 81–87. <https://doi.org/10.1016/j.humov.2018.03.016>
27. Izovska, J., Mikic, M., Dragijsky, M., Zahalka, F., Bujnovsky, D., & Hank, M. (2019). Pre-Season Bilateral Strength Asymmetries of Professional Soccer Players and Relationship with Non-Contact Injury of Lower Limb in the Season. *Sport Mont*, 17(2). <https://doi.org/10.26773/smj.190619>
28. Carey, D. P., Smith, G., Smith, D. T., Shepherd, J. W., Skriver, J., Ord, L., & Rutland, A. (2001). Footedness in world soccer: An analysis of France '98. *Journal of Sports Sciences*, 19(11), 855–864. <https://doi.org/10.1080/026404101753113804>
29. DeLang, M. D., Salamh, P. A., Farooq, A., Tabben, M., Whiteley, R., van Dyk, N., & Chamari, K. (2021). The dominant leg is more likely to get injured in soccer players: Systematic review and meta-analysis. *Biology of Sport*, 38(3), 397–435. <https://doi.org/10.5114/biolSport.2021.100265>
30. Häggglund, M., Waldén, M., & Ekstrand, J. (2012). Risk factors for lower extremity muscle injury in professional soccer. *The American Journal of Sports Medicine*, 41(2), 327–335. <https://doi.org/10.1177/0363546512470634>
31. Serner, A., Tol, J. L., Jomaah, N., Weir, A., Whiteley, R., Thorborg, K., Robinson, M., & Hölmich, P. (2015). Diagnosis of acute groin injuries: A prospective study of 110 athletes. *The American Journal of Sports Medicine*, 43(8), 1857–1864. <https://doi.org/10.1177/0363546515585123>
32. Mancino, F., Gabr, A., Plastow, R., & Haddad, F. S. (2023). Anterior cruciate ligament injuries in female athletes. *The Bone & Joint Journal*, 105-B(10), 1033–1037. <https://doi.org/10.1302/0301-620X.105B10.BJJ-2023-0881.R1>
33. Hewett, T. E., Myer, G. D., & Zazulak, B. T. (2008). Hamstrings to quadriceps peak torque ratios diverge between sexes with increasing isokinetic angular velocity. *Journal of Science and Medicine in Sport*, 11(5), 452–459. <https://doi.org/10.1016/j.jsams.2007.04.009>
34. Knapik, J. J., Bauman, C. L., Jones, B. H., Harris, J. M., & Vaughan, L. (1991). Preseason strength and flexibility imbalances associated with athletic injuries in female collegiate athletes. *The American Journal of Sports Medicine*, 19(1), 76–81. <https://doi.org/10.1177/036354659101900113>

-
35. Behan, F. P., Maden-Wilkinson, T. M., Pain, M. T. G., & Folland, J. P. (2018). Sex differences in muscle morphology of the knee flexors and knee extensors. *PloS One*, 13(1), e0190903. <https://doi.org/10.1371/journal.pone.0190903>
 36. De Ste Croix, M., ElNagar, Y. O., Iga, J., Ayala, F., & James, D. (2017b). The impact of joint angle and movement velocity on sex differences in the functional hamstring/quadriceps ratio. *The Knee*, 24(4), 745–750. <https://doi.org/10.1016/j.knee.2017.03.012>
 37. O'Sullivan, L., & Tanaka, M. J. (2021). Sex-based differences in hamstring injury risk factors. *Journal of Women's Sports Medicine*, 1(1), 20-29. <https://doi.org/10.53646/jwsm.v1i1.8>
 38. Śliwowski, R., Paillard, T., Bojkowski, Ł., Dudziński, W., Patek, M., & Marynowicz, J. (2024). Intra- and inter-limb strength imbalance and asymmetry in soccer: A comparison of elite senior and junior players. *PLoS ONE*, 19(4), e0302474. <https://doi.org/10.1371/journal.pone.0302474>